

# String pair production in non homogeneous backgrounds

*Stefano Bolognesi*

**Università di Pisa and INFN**

arXiv:1601.04785 in collaboration with E. Rabinovici and G.Tallarita

**CAQCD16**

# Plan of the talk

Introduction to pair production, worldline formalism and non homogeneous backgrounds in field theory



# Plan of the talk

Introduction to pair production, worldline formalism and non homogeneous backgrounds in field theory

String theory examples of pair production which are treatable with the “worldsheet instanton” technique:

- 1) String suspended between D-branes
- 2) Holographic Schwinger effect

# Plan of the talk

Introduction to pair production, worldline formalism and non homogeneous backgrounds in field theory


String theory examples of pair production which are treatable with the “*worldsheet instanton*” technique:

- 1) String suspended between D-branes
- 2) Holographic Schwinger effect


Effects of non-homogeneous backgrounds on string production

# Pair production

Schwinger effect is non-perturbative pair production of  $q$ - $q^*$  due to electric background




$$\text{Im } \mathcal{L}[E] = \frac{e^2 E^2}{16\pi^3} \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^2} \exp \left[ -\frac{m^2 \pi n}{eE} \right]$$




# Pair production

Schwinger effect is non-perturbative pair production of  $q$ - $q^*$  due to electric background



This effect becomes significant at

$$E \simeq m^2/e$$



# Worldline instanton

One searches for stationary solution of the Euclidean worldline action

$$S_E = m \int d\tau \sqrt{\dot{x}^\mu \dot{x}^\mu} + iq \int d\tau \dot{x}^\mu A^\mu$$

Same trajectories of a charged particle moving in a background magnetic field

$$\frac{m\ddot{x}^\mu}{2\sqrt{\dot{x}^\mu \dot{x}^\mu}} = iqF_{\mu\nu}\dot{x}^\nu$$

$$\dot{x}^\mu \dot{x}^\mu = \text{const} = L^2$$


# Worldline instanton

For a constant background the solution is given by a circular trajectory

$$x_3(\tau) = \frac{m}{qE} \cos(2\pi\tau) \quad x_4(\tau) = \frac{m}{qE} \sin(2\pi\tau)$$

with action consistent with Schwinger formula


$$S_E = \frac{\pi m^2}{qE}$$






# Pair production in *QFT*

Solutions are “extremal” circular orbits with radius  $R$



# Pair production in *QFT*



They describe the tunneling in Minkowsky

$$S_E = \frac{\pi m^2}{qE}$$

$$P \propto \exp(-S_E)$$

# Non homogeneous background

Schwinger effect becomes significant at  $E \simeq m^2/e$

Non homogeneous backgrounds can lower this value significantly!

Direct observation of the Schwinger effect may be possible in the near future with the use of strong laser pulses


*Sauter  
Brezin-Itzykson 70*

...

# Time-dependent pulse

The simplest case is a single pulse of electric field dependent on time only

$$E^3(t) = \frac{E}{\cosh^2(\omega t)}$$




# Time-dependent pulse

The simplest case is a single pulse of electric field dependent on time only

$$E^3(t) = \frac{E}{\cosh^2(\omega t)}$$

The Euclidean corresponding field is

$$F_{34} = \frac{-iE}{\cos^2(\omega x_4)}$$



We already see that there should be enhancement of pair production if the instanton is finite!


# Time-dependent pulse

The exact solution is:



$$x_3(\tau) = \frac{1}{\omega} \frac{1}{\sqrt{1 + \gamma^2}} \operatorname{arcsinh}(\gamma \cos(2\pi\tau))$$

$$x_4(\tau) = \frac{1}{\omega} \arcsin\left(\frac{\gamma}{\sqrt{1 + \gamma^2}} \sin(2\pi\tau)\right)$$

$$\gamma = \frac{m\omega}{qE}$$



# Time-dependent pulse




# Time-dependent oscillation

Another important case is an oscillating electric field with fixed frequency

$$E^3(t) = E \cos(\omega t)$$

The Euclidean corresponding field is



$$F_{34} = -iE \cosh(\omega x_4)$$






# Time-dependent oscillation

The exact solution is:



# Summary of Results


Worldline action comparison:



Both time dependent actions are lower than the “locally constant” approximation. This indicate enhancement of pair production


# Summary of Results

Worldline action comparison:




Continuous interpolation  
between “tunnelling” and  
“photon absorption”


# Space-dependent pulse



# Space-dependent oscillation



# Summary of Results



For space dependent backgrounds is just the opposite

# Pair production in string theory

In string theory there is a critical value where the barrier for pair production disappears and the vacuum is “catastrophically” unstable

$$eE_{cr} = \frac{1}{2\pi\alpha'}$$

*Fradkin-Tseytlin 85*

*Burgess 87*

*Bachas-Porrati 92*

...


Not much is known about string pair production in non homogeneous backgrounds

# String suspended between two D-branes

We look for a case where pair production can be studied with the “worldsheet instanton” technique



# String suspended between two D-branes



We look for a case where pair production can be studied with the “worldsheet instanton” technique

$$m = Td$$

$$S = T \int d\sigma d\tau \sqrt{\det g_2(\sigma, \tau)} + iq \int_{\text{boundary}} dX^\mu A_\mu$$

# Minimal surface solution

For constant background, and so circular symmetry:

$$S_E = T \int_{-d/2}^{d/2} dz 2\pi r(z) \sqrt{1 + r'(z)^2} - 2qE\pi R^2$$

# Minimal surface solution


For constant background, and so circular symmetry:

$$S_E = T \int_{-d/2}^{d/2} dz 2\pi r(z) \sqrt{1 + r'(z)^2} - 2qE\pi R^2$$

The solutions are the well known “catenaries” minimal surface




$$r(z) = \frac{1}{c} \cosh(cz)$$

$$R = \frac{1}{c} \cosh(cd/2)$$




# String suspended between two D-branes

For  $R$  sufficiently large there are two solutions: “thick neck” and “thin neck”



# String suspended between two D-branes

For  $R$  sufficiently large there are two solutions: “thick neck” and “thin neck”




Critical field is as expected:  $qE_{\text{cr}} = T$

# Time-dependent setup

We work in cylindrical coordinates  $(z, r, \theta)$

and have to find a function of two variables  $r(z, \theta)$



$$S_E = T \int_{-d/2}^{d/2} dz \int_0^{2\pi} d\theta \sqrt{r^2(1 + (\partial_z r)^2) + (\partial_\theta r)^2} - iq \int_0^{2\pi} d\theta (A_\theta + A_r \partial_\theta r)$$

# Time-dependent setup

The boundary term is given by the specific choice of background electric field

$$A_\theta = -iE f(r \sin \theta) r \sin \theta$$




$$A_r = iE f(r \sin \theta) \cos \theta$$

$$f(x_4) = \frac{\tan(\omega x_4)}{\omega} \quad \text{pulse}$$




$$f(x_4) = \frac{\sinh(\omega x_4)}{\omega} \quad \text{oscillating}$$

We use Fourier angular decomposition and Chebyshev polynomials in  $z$

# Solution for small frequency case






Pulse  
with  $\omega = 0.1$








# Solution for small frequency case



Oscillation  
with  $\omega = 0.1$



# Solution for higher frequency

String effects are expected to be important when  $E$  is big enough

$$qE \simeq T.$$




Non-homogeneity effects are expected to be important when  $E$  is small enough

$$qE < m\omega$$

So we need to increase  $\omega$  to see effects that are both “stringy” and related to non-homogeneity




# Solution for higher frequency

Pulse  
with  $\omega = 0.3$



# Solution for higher frequency




Pulse  
with  $\omega = 0.3$




“Double” enhancement of pair production

# Solution for higher frequency

Oscillation  
with  $\omega = 0.4$



# Small E string effects (?)




As  $E \rightarrow 0$  the area inside the loop goes to zero and becomes smaller than worldsheet area




$$\frac{TqE_{\text{low}}}{\pi m^2 \omega^2} \log \left( \frac{2m\omega}{qE_{\text{low}}} \right) \simeq 1$$



# Small E string effects (?)



Pulse




As  $E \rightarrow 0$  the area inside the loop goes to zero and becomes smaller than worldsheet area



$$\frac{TqE_{\text{low}}}{\pi m^2 \omega^2} \log \left( \frac{2m\omega}{qE_{\text{low}}} \right) \simeq 1$$

$$E_{\text{low}} \simeq 0.0026$$


# Space dependent pulse



Pulse  
with  $k = 0.2$




$E = 0.3$





$E = 0.9$




# Space dependent oscillation



Oscillation  
with  $k = 0.2$



$E = 0.3$



$E = 0.9$


# Holographic Schwinger effect

We consider N=4 SYM in the Coulomb phase

N D3

1 D3


$$SU(N+1) \rightarrow SU(N) \times U(1)$$



This is the U(1) of the Schwinger effect

# Holographic Schwinger effect

We consider  $N=4$  SYM in the Coulomb phase



# Holographic Schwinger effect

The dictionary is the usual

$$g_s = g^2/4\pi \quad L^2/l_s^2 = \sqrt{\lambda}/2\pi$$

plus the relation for the W mass

$$\frac{1}{l_s^2} \int_0^{r_0} \sqrt{-\det h_{ab}} = \frac{L^2 r_0}{l_s^2} = \frac{\sqrt{\lambda} r_0}{2\pi}$$

$$m = \frac{TL^2}{z_0}$$



# *Holographic setting*

The DBI action predicts the string to break down at “local” string scale

This is translated into a genuine QFT scale

$$S_{DBI} = \frac{1}{g_s l_s^4} \int d^4 x' \sqrt{-\det(\eta_{\mu\nu} - l_s^2 F_{\mu\nu, loc})}$$


This is the value where the barrier for the pair production drops to zero in the QFT!

$$E_{cr} = \frac{r_0^2 L^2}{l_s^2} = \frac{2\pi m^2}{\sqrt{\lambda}}$$

# Holographic worldsheet instanton

In holography, the instanton is a Wilson loop attached to the D3 brane

$$ds^2 = \frac{L^2}{z^2} (dz^2 + dx_\mu dx^\mu)$$




# Holographic worldsheet instanton

$$S_E = T \int_0^R d\rho 2\pi\rho \left( \frac{L}{z(\rho)} \right)^2 \sqrt{1 + z'(\rho)^2} - qE\pi R^2$$

Minimal surface is a spherical cap  $z(\rho) = \sqrt{R_{\text{UV}}^2 - \rho^2}$

Boundary condition  $T \cos(\gamma) = qF_{\text{loc}}$



# Holographic worldsheet instanton

The action  $S_E = \frac{\pi}{qE z_0^2} \left( \left( \frac{TL^2}{qE z_0^2} \right) - 1 \right)^2$  vanishes when the spherical cap is tangent to the D-brane

$$qE_{\text{cr}} = \frac{TL^2}{z_0^2}$$




# Holographic worldsheet instanton

Tunneling barrier exists only below a certain critical field

$$S_E = \frac{\mathcal{T} \pi L^2 R^2}{z_0^2} - q F \pi R^2 + \dots$$

$$q F_{cr} = \frac{\mathcal{T} L^2}{z_0^2}$$



# Time-dependence setup

As before:


we work in cylindrical coordinates  $(z, r, \theta)$

and have to find a function of two variables  $r(z, \theta)$

$$S_E = T \int_{z_0}^{Z_0} dz \int_0^{2\pi} d\theta \frac{L^2}{z^2} \sqrt{r^2(1 + (\partial_z r)^2) + (\partial_\theta r)^2} - iq \int_0^{2\pi} d\theta (A_\theta + A_r \partial_\theta r)$$

# Time-dependence setup

Boundary conditions are:






$r(Z_0, \theta) = 0,$        $r'(Z_0, \theta) = \infty$


$$T \frac{L^2}{z_0^2} \frac{r^2 \partial_z r}{\sqrt{r^2 (1 + (\partial_z r)^2) + (\partial_\theta r)^2}} = q E f' (r - \partial_\theta r \sin \theta \cos \theta)$$

# Some solutions

Pulse  
with  $\omega = 0.15$






$E = 0.4$




$E = 0.9$

# Some solutions

Oscillation  
with  $\omega = 0.15$



$E = 0.4$




$E = 0.9$

# Confining background


Hard-wall geometry:

Confining string


$$T_c = T \frac{L^2}{z_{\text{IR}}^2}$$



# Confining background



$$E > E'$$





$$E < E'$$

At the field  $qE' = \frac{TL^2}{z_0 z_{\text{IR}}}$

we have a transition in the type of worldsheet solution:

# Low critical field in confing background



$$qE'_{cr} = \frac{TL^2}{z_{IR}^2} = T_c$$



# Low critical field in confining background

The interpretation of the “low” critical field in Minkowski:

$$V(R) = TL^2 \left( \frac{1}{z_0} - \frac{1}{z_{\text{IR}}} \right) + T_c R - qER$$

When  $qE > T_c$ , the electric field is not strong enough to overcome confinement

# Low critical field and time-dependence

The critical low field  $E'_{\text{cr}}$  is in general modified by the frequency  $\omega$ .


We show a clear example in which it vanishes!

# Low critical field and time-dependence

Maximum holographic  $z$  for low  $E$ :


Unlike the constant background case,  
 $z_{\max}$  is bounded from above:

$$z_{\max} = z_0 + \frac{c\pi}{2\omega} \simeq \frac{2.62}{\omega}$$



# Low critical field and time-dependence

Maximum holographic  $z$  for low  $E$ :



If the confinement scale is so that

$$z_{\text{IR}} > z_{\text{max}}$$

The Euclidean worldsheet is always contained inside the physical space.

Pair production happen irrespectively on how small is  $E$ .

Photons energy is enough to produce glueballs and charged particles are intermediary states

# Conclusion

- We studied two cases of string pair production with the technique of Euclidean worldsheet instanton
- With this technique we can define the problem on non homogeneous backgrounds and solve PDE numerically
- String nature and non homogeneity work together when  $\omega$  is big enough; for time dependent backgrounds they enhance even further the pair production
- Pair production in confining background is highly modified by the non-homogeneity. In particular the low critical field can disappear